



SUMMARY

- Angus HeiferSELECT is a genomic selection tool to help inform the selection of Angus replacement females (87.5% Angus content or greater) in a commercial beef breeding operation.
- The 2nd generation of Angus HeiferSELECT was commercialised in November 2021 following 18 months of close collaboration between Angus Australia and CSIRO (Australia's National Science Agency).
- The fertility genetic prediction, Heifer Conception, was added to the Angus HeiferSELECT suite in August 2023, based on research conducted in partnership with CSIRO.
- Angus HeiferSELECT provides genetic predictions for fourteen (14) maternal, fertility, growth, feed intake, carcass and resilience traits.
- The reference population (Angus animals with genotypes and phenotypes) that underpins the Angus HeiferSELECT genetic prediction is based on the comprehensive data from the Angus Australia database, including hard-to-measure traits (carcass, feed intake, immune competence) from Angus Australia's reference population program, known as the Angus Sire Benchmarking program (ASBP).
- Angus HeiferSELECT provides three overall selection Indexes being Cow-Calf Value (CCV), Feedlot and Carcass Value (FCV) and Total Breeding Value (TBV) which have been developed in collaboration with AbacusBio (a global leader in the development of livestock selection indexes) and are based on the latest selection index economic modelling and current day production parameters.
- An addition to this product is Angus BreedCHECK, which is a genomic (DNA) based system that estimates breed composition (from 11 breeds), with a particular focus on Angus content
- An important feature of Angus HeiferSELECT is the comprehensive validation that has been undertaken to ensure it is an effective selection tool, particularly for Australian Angus in Australian production systems

CONTENTS

1. BACKGROUND

2. FEATURES

2.1. Genetic Predictions

2.1.1. Traits

2.1.2. Model and reference Population

2.2. Overall Values

2.2.1. Cow-Calf Value (CCV)

2.2.2. Feedlot and Carcase Value (FCV)

2.2.3. Total Breeding Value (TBV)

2.3. Angus Breedcheck

2.4 Sire Assignment

2.5 Add-on BVDV testing

3. GENETICS EFFECTS TABLE

4. VALIDATION

1. BACKGROUND

Angus HeiferSELECT is a genomic selection tool to help inform the selection of Angus replacement females (87.5% Angus content or greater) in a commercial beef breeding operation.

Angus HeiferSELECT provides:

- Genetic predictions for fourteen (14) maternal, fertility, growth, feed intake, carcase and resilience traits
- Genetic prediction for cow-calf value, feedlot-carcase value and total breeding value (with star rating)
- Angus BreedCHECK – genomic breed composition prediction
- Sire assignment – Possible sires must be registered with Angus Australia and have genomic profiles available

Angus HeiferSELECT complements other sources of information that may be used in commercial replacement heifer selection, such as phenotypic evaluation, age, weight and pedigree information, and provides valuable insight into the genetic potential of heifers, particularly for traits that are otherwise difficult, time consuming or expensive to measure using traditional methods.

The first generation of Angus HeiferSELECT was commercialised in 2017. The 2nd generation of Angus HeiferSELECT was commercialised in November 2021 following 18 months of close collaboration between Angus Australia and CSIRO (Australia's National Science Agency).

Further research conducted in partnership with CSIRO culminated in the release of a fertility genetic prediction, Heifer Conception, being released in August 2023.

Angus HeiferSELECT is delivered to Angus Australia members in collaboration with Zoetis and Neogen Australasia.



zoetis



NEOGEN
Australasia

2. FEATURES

2.1. Genetic Predictions

2.1.1. Traits

Angus HeiferSELECT provides genetic predictions for fourteen (14) maternal, fertility, growth, feed intake, carcase and resilience traits. The genetic predictions are reporting using an intuitive 0 – 100 scoring system, with a score of 50 representing the average genetic merit of commercial Angus heifers tested with the Angus HeiferSELECT product.

Higher values identify females carrying genetics that will produce “more” of a trait, which may or may not be preferred, subject to your breeding objective. For example, a female with a Yearling Weight genetic prediction of 80 would be expected to produce progeny that are heavier at 13-14 months of age than a female with a Yearling Weight genetic prediction of 30, all other things being equal. Similarly, higher HC genetic predictions indicate the animal is expected to produce daughters that are more likely to conceive and conceive earlier as heifers (older foetal age at pregnancy diagnosis).

A description of each of the traits by group are listed in table 1.

2.1.2. Model and Reference Population

The model used for calculating the Angus HeiferSELECT genetic predictions (i.e. genomic breeding values) is based on applying SNP effects to 45,364 SNPs in a standard imputed genomic profile for each of the 14 traits, for each heifer. The SNP effects resulting from univariate analyses of each trait.

The reference population (Angus animals with genotypes and phenotypes) that underpins the Angus HeiferSELECT genetic predictions is based on the comprehensive data from the Angus Australia database, including hard-to-measure traits (carcase, feed intake, immune competence) from Angus Australia's reference population program, known as the Angus Sire Benchmarking program (ASBP) (Table 2). All animals in the reference population are straight-bred Angus from Australian production systems.

Table 1 - Description of 14 Angus HeiferSELECT Genetic Predictions

TRAIT		DESCRIPTION
COW-CALF TRAITS	Calving Ease	Higher Calving Ease (CE) genetic predictions indicate the animal is expected to experience lower birth weight and fewer calving difficulties as a 2 year old heifer (i.e. greater calving ease).
	Weaning Weight	Higher Weaning Weight (WW) genetic predictions indicate the animal is expected to produce progeny with heavier live weights at 200 days of age, due to superior growth potential.
	Milk	Higher Milk genetic predictions indicate the animal is expected to produce progeny with heavier live weights at 200 days of age, due to superior maternal attributes (i.e. more milk).
	Yearling Weight	Higher Yearling Weight (YW) genetic predictions indicate the animal is expected to produce progeny with heavier live weights at 400 days of age.
	Mature Cow Weight	Higher Mature Cow Weight (MCW) genetic predictions indicate the animal is expected to have a heavier weight at 3.5 years of age, and produce female progeny that are heavier. Heavier mature weights are associated with higher feed and maintenance costs, but conversely higher returns for cull cows.
	Heifer Conception	Higher Heifer Conception (HC) genetic predictions indicate the animal is expected to produce daughters that are more likely to conceive and conceive earlier as heifers (older foetal age at pregnancy diagnosis), than those with lower values.
FEEDLOT TRAITS	Average Daily Gain	Higher Average Daily Gain (ADG) genetic predictions indicate the animal is expected to produce progeny with higher rates of weight gain during feedlot finishing, due to superior growth potential.
	Daily Feed Intake	Higher Daily Feed Intake (DFI) genetic predictions indicate the animal is expected to produce progeny that eat more during feedlot finishing, and may be considered less efficient than lower
CARCASE TRAITS	Carcase Weight	Higher Carcase Weight (CW) genetic predictions indicate the animal is expected to produce progeny with heavier carcase weights.
	Eye Muscle Area	Higher Eye Muscle Area (EMA) genetic predictions indicate the animal is expected to produce progeny with more muscle and larger eye muscle area.
	Rib Fat	Higher Rib Fat (RIB) genetic predictions indicate the animal is expected to produce progeny with greater fat depth.
	MSA Marbling	Higher MSA Marbling (MBL) genetic predictions indicate the animal is expected to produce progeny with higher marbling scores and more intramuscular fat
	Ossification	Higher Ossification (OSS) genetic predictions indicate the animal is expected to produce progeny with higher levels of ossification, or physiological maturity, in the carcase which is antagonistic to eating quality.
RESILIENCE	ImmuneDEX	Higher ImmuneDEX (IMM) genetic predictions indicate the animal is expected to produce progeny with higher levels of general disease resilience, as measured by cell-mediated and antibody mediated immune response.

Table 2. Summary of the Angus Australia Reference Population that Underpins Angus HeiferSELECT (Generation 2)

TRAIT	N	MEAN	SD	Vg	h2
Birth Weight (kg)	50,504	37.2	5.02	7.31	44
Weaning Weight (kg)	46,106	249.2	50.57	135.84	25
Yearling Weight (kg)	36,745	416.4	79.5	305.67	30
Mature Cow Weight (kg)	2,683	550.1	77.1	973.08	47
Heifer Conception (weeks)	6,070	13	7.4	11.74	32
Average Daily Gain (kg/d)	3,993	1.6	0.339	0.02	27
Daily Feed Intake (kg/d)	3,993	14.4	2.078	0.88	34
Carcase Weight (kg)	3,764	427.1	66.85	495.19	49
Eye Muscle Area (cm ²)	3,755	89.4	10.87	31.16	44
MSA Marbling Score	3,762	492.3	125.88	4008.64	40
Ossification Score	3,761	147.3	18.63	58.14	32
Rib Fat (mm)	3,741	16.8	6.15	6.51	29
ImmuneDEX (Index)	3,715	0.005	1.137	0.25	26

2.2. Overall Values

Angus HeiferSELECT provides three overall values based on economic selection index modelling being Cow-Calf Value (CCV), Feedlot and Carcase Value (FCV) and Total Breeding Value (TBV).

The overall values have been developed in collaboration with AbacusBio (a global leader in the development of livestock selection indexes) and are based on the latest selection index economic modelling and current day production parameters. This is further outlined in a conference paper "*Indexes supporting genomic tools for selecting commercial Angus heifer replacements and identifying steers for long-fed programmes in Australia*" delivered at the 24th Association for the Advancement of Animal Breeding and Genetics (AAABG) conference.

Reference: Quinton C. D. et al (2021). Indexes Supporting Genomic Tools For Selecting Commercial Angus Heifer Replacements And Identifying Steers For Long-Fed Programmes In Australia. Assoc. Advmt. Anim. Breed. Genet: 24: 308-311. <http://www.aaabg.org/aaabghome/AAABG24papers/77Quinton24308.pdf>

2.2.1. Cow-Calf Value (CCV)

Cow-Calf Value (CCV) estimates the genetic differences between animals in net profitability in a typical commercial Angus self-replacing herd, focusing on the traits related to the cow-calf production system. The Cow-Calf Value assists in making "balanced" selection decisions, taking into account the relevant calving ease, growth and maternal attributes to identify animals that are most suitable for use within a particular commercial enterprise. Higher Cow-Calf value genetic predictions identifies animals that will improve overall profitability in the majority of commercial systems selecting Angus females.

The Cow-Calf Value is a sub-index (i.e. component) of the Total Breeding Value.



Figure 1 shows the traits that are considered in the CCV, and how much they contribute to the overall balance of the value. The larger the segment, the greater the impact on the selection index. In the CCV, there is a focus on increasing weaning weight and milk, while maintaining calving ease and mature cow weight.

Figure 1. Trait Emphasis for the Cow-Calf Value (CCV)





Figure 2 shows the selection advantage if heifers are selected using the CCV. The selection advantage is calculated by ranking a group of Angus heifers on CCV, and comparing the average genetic predictions of the heifers in the top 30% with the average genetic predictions of all heifers available for selection.

Figure 2. Selection Advantage for Cow-Calf Value



2.2.2. Feedlot and Carcase Value (FCV)

Feedlot-Carcass Value (FCV) estimates the genetic differences between animals in net profitability in a typical commercial Angus self-replacing herd, focusing on the traits related to feedlot and processor performance, along with consumer expectations. The Feedlot-Carcass Value assists in making "balanced" selection decisions, taking into account the relevant growth, feed intake and carcass attributes to identify animals that are most suitable for use within a particular commercial enterprise. Higher Feedlot-Carcass Value genetic predictions identifies animals that will improve overall profitability in the majority of commercial systems selecting Angus females with progeny entering grain finishing supply chain.

Figure 3 shows the traits that are considered in the FCV, and how much they contribute to the overall balance of the value. The larger the segment, the greater the impact on the selection index. In the FCV, there is a focus on increasing growth resulting in heavier carcass weights and MSA marbling, while maintaining rib fat.

Figure 4 shows the selection advantage if heifers are selected using the FCV. The selection advantage is calculated by ranking a group of Angus heifers on FCV, and comparing the average genetic predictions of the heifers in the top 30% with the average genetic predictions of all heifers available for selection.

The Feedlot-Carcass Value is a sub-index (i.e. component) of the Total Breeding Value

Figure 3. Trait Emphasis for the Feedlot-Carcass Value





Figure 4. Selection Advantage for Feedlot-Carcass Value



2.2.3. Total Breeding Value (TBV)

Total Breeding Value (TBV) estimates the genetic differences between animals in net profitability in a typical commercial Angus self-replacing herd with progeny entering the grain finishing supply chain. The Total Breeding Value assists in making "balanced" selection decisions, taking into account the relevant calving ease, growth, feed intake and carcass attributes to identify animals that are most suitable for use within a particular commercial enterprise. Higher Total breeding value genetic predictions identifies animals that will improve overall profitability in the majority of commercial systems selecting Angus females.

The Cow-Calf Value and Feedlot-Carcass value are sub-indices (i.e. components) of the Total Breeding Value.

Figure 5. Trait Emphasis for the Total Breeding Value





Figure 5 shows the traits that are considered in the TBV, and how much they contribute to the overall balance of the value. The larger the segment, the greater the impact on the selection index. In the TBV, there is a focus on increasing growth resulting in heavier carcass weights and more MSA marbling, while maintaining calving ease and mature cow weight.

Figure 6 shows the selection advantage if heifers are selected using the TBV. The selection advantage is calculated by ranking a group of Angus heifers on TBV, and comparing the average genetic predictions of the heifers in the top 30% with the average genetic predictions of all heifers available for selection.

Figure 6. Selection Advantage for the Total Breeding Value



2.3. Angus Breedcheck

Angus BreedCHECK is a genomic (DNA) based system that estimates breed composition (from 11 breeds), with a particular focus on Angus content.

Heifers that are 87.5% (or 7/8th) or greater Angus content receive the Angus BreedCHECK tick. Heifers that are below 87.5% Angus content are flagged with an Angus BreedCHECK cross and provided with additional information to further understand their breed background. This includes the percentage (%) content value

estimate for:

- Angus
- Non-Angus
- British (including Angus, Hereford, Shorthorn and Murray Grey)
- Indicus (including Brahman and Santa Gertrudis)
- European (including Charolais, Simmental and Limousin)

· Other (including Holstein and Wagyu)

Angus HeiferSELECT genetic predictions are provided on all animals that are greater than 50% Angus content. Caution should be applied when using the Genetic Predictions for animals less than 87.5% Angus as the associated reference population is based on straight bred Angus animals.

The software underpinning Angus BreedCHECK was developed in collaboration with the CSIRO (Australia's National Science Agency). This is based on an allele frequency approach, with effects being applied to 45,364 SNPs in a standard imputed genomic profile for each of the 11 breeds, for each heifer.

There are 1,850 animals in the reference population (animals with known breed and genotypes) that underpins Angus BreedCHECK as listed in table 3.



Table 3. Summary of the Reference Population that Underpins Angus BreedCHECK

BREED	REFERENCE
Angus	868
Brahman	330
Charolais	71
Hereford	111
Holstein	144
Limousin	53
Murray Grey	62
Santa Gertrudis	53
Shorthorn	88
Simmental	27
Wagyu	43
Total	1,850

2.4. Sire Assignment

DNA based sire assignment is included as a feature of Angus HeiferSELECT.

An essential prerequisite for sire determination, is that candidate sires must be registered with Angus Australia and have genomic profile available.

In many cases, a genomic (DNA) profile will have previously been recorded with Angus Australia for sires (by the breeder or a previous owner) and can be used to conduct the DNA sire identification component of the Angus HeiferSELECT testing. Details of whether a DNA profile is stored for each registered sire can be viewed on the Angus Database Search facility on the Angus Australia website.

If a genomic profile has not previously been recorded, a DNA sample can be collected for the sire and request a genomic profile from Angus Australia. DNA test request forms are available from the Angus Australia website.

2.5. Add-on BVDV Testing

Bovine viral diarrhoea virus (BVDV) in cattle is a complex disease that is caused by bovine pestivirus. Bovine pestivirus and its resulting diseases have several interchangeable names including bovine viral diarrhoea (BVD), pestivirus and bovine pestivirus. Mucosal disease is the fatal disease that develops in animals persistently infected with BVDV.

BVDV can reduce herd reproductive rates through infertility or abortion, as well as reduce animal's immunity to a range of other diseases such as bovine respiratory disease (BRD). BVDV is common in Australia and Meat and Livestock Australia (MLA) have identified BVDV as the second most costly disease to Australian cattle herds, after cattle tick infection, and the most important cattle disease southern cattle herds. It is estimated to have an economic impact of \$114 million per year.

BVDV testing can be conducted in association with Angus HeiferSELECT as an optional add-on from both Zoetis (TSU samples only) and Neogen (TSU and tail hair samples). This testing facilitates the identification of animals persistently infected with BVDV.

If BVDV testing is required, this should be nominated on the order form. Angus HeiferSELECT and BVDV testing can be conducted from the same DNA sample.



Table 4. Angus HeiferSELECT Genetic Effects Table

CE (KG)	WW (KG)	YW (kg)	Milk (kg)	MCW (kg)	HC (weeks)	ADG (kg/day)	DFI (kg/day)	CW (kg)	EMA (cm ²)	RIB (mm)	MBL (score)	OSS (score)	CCV (\$)	FCV (\$)	TBV (\$)
-0.67	2.1	3.5	1.3	6.7	0.6	0.03	0.23	6.1	1.4	0.5	17.9	1.6	12.9	36.3	39.7

3. GENETICS EFFECTS TABLE

The Angus HeiferSELECT genetic effects table (Table 4.) enables you to assess genetic differences between heifers tested with Angus HeiferSELECT in real terms by providing the predicted genetic difference per ten (10) unit change in Angus HeiferSELECT value.

The genetic effects table can be used to predict the difference in progeny performance between two females. For example, a female with a Carcase Weight genetic prediction of 80 would be expected to produce progeny that are, on average, 15.25 kg heavier at around 2 years of age, compared to a female with a Carcase Weight genetic prediction of 30, all other things being equal.

[Working: 6.1 kg genetic difference per 10 unit HeiferSELECT value, 30.5 kg (6.1 x 5) genetic difference per 50 unit HeiferSELECT value, 15.25 kg (30.5 / 2) progeny difference (as females only contribute half of the genetics to the progeny, with the remainder coming from the sire to which they are joined)]

Also note:

- Calving Ease is presented in birth weight units of kgs, with higher CE genetic predictions resulting in lower birth weight, a major contributor to overall calving ease.
- The CCV, FCV and TBV are based of economic modelling underpinned by current day parameters therefore Australian dollar (\$) units apply.

4. VALIDATION

An important feature of Angus HeiferSELECT is the comprehensive validation that has been undertaken to ensure it is an effective selection tool, particularly for Australian Angus in Australian production systems.

The validation includes a combination of peer reviewed journal papers and internal validation.

Three peer reviewed papers, focusing on the carcass trait genetic predictions, show Angus HeiferSELECT (and the related product Angus SteerSELECT) can predict differences in phenotype and performance. The papers are:

- *“Development of Angus SteerSELECT: a genomic-based tool to identify performance differences of Australian Angus steers during feedlot finishing: Phase 1 validation”* with conclusion: Genomic prediction equations can predict differences in carcass weight, marbling score, ossification score and carcass value in both short-fed and long-fed Australian Angus.

Reference: Hine B. C. et al (2021). Development of Angus SteerSELECT: a genomic-based tool to identify performance differences of Australian Angus steers during feedlot finishing: Phase 1 validation. *Animal Production Science* 61, 1884-1892. <https://doi.org/10.1071/AN21051>

- *“Bias, dispersion, and accuracy of genomic predictions for feedlot and carcass traits in Australian Angus steers”* with conclusion: Estimates of h² and GEBV quality metrics suggest a potential for accurate genomic selection of Australian Angus for feedlot performance and carcass traits.

Reference: Alexandre, P.A. et al (2021) Dispersion, and accuracy of genomic predictions for feedlot and carcass traits in Australian Angus steers. *Genet Sel Evol* 53, 77. <https://doi.org/10.1186/s12711-021-00673-8>

- *“Development of female fertility indicator traits for the Angus HeiferSELECT genomic tool”* with conclusion: Selection based on Heifer Conception (PREG) can result in tangible gains in female fertility in Angus females in commercial enterprises through the use of Angus HeiferSELECT.

Reference: Alexandre, P.A. et al (2023) Development of female fertility indicator traits for the Angus HeiferSELECT genomic tool. *Proc. Assoc. Advmt. Anim. Breed. Genet.* 25: 67 - 70.

Internal Validation (unpublished) has also shown that Angus HeiferSELECT genetic predictions for calving ease, weaning weight, yearling weight and mature cow weight can predict phenotype and performance differences, both on the heifers themselves (Table 5) and for their progeny (Table 6). For example, for yearling weight (YW), the difference between the top and bottom quartiles for self-performance and progeny performance is 19.55 kg and 9.55 kg respectively.

Note:

- Calving Ease (CE) is presented in birth weight units of kgs, with higher CE genetic predictions resulting in lower birth weight, a major contributor to overall calving ease.
- The phenotypes presented against each quartile are values adjusted for age, age of dam and contemporary group.




Table 5. Quartile analyses for HeiferSELECT Genetic Predictions for CE, WW, YW and MCW based on self-performance

	N	CE (KG)	N	WW (KG)	N	YW (kg)	N	MCW (kg)
Q1	30,709	1.84	27,544	8.59	21,772	12.20	1,998	33.02
Q2	30,709	0.56	27,545	3.75	21,773	4.80	1,998	9.26
Q3	30,709	-0.31	27,545	0.73	21,773	-0.24	1,998	-3.52
Q4	30,710	-1.69	27,545	-4.39	21,773	-7.34	1,998	-28.11
Q1Q4 Diff		3.53		12.98		19.55		61.13

Table 6. Quartile analyses for HeiferSELECT Genetic Predictions for CE, WW, YW and MCW based on progeny performance

	N	CE (KG)	N	WW (KG)	N	YW (kg)	N	MCW (kg)
Q1	8,124	0.86	6,570	3.59	5,364	5.46	1,064	12.92
Q2	8,125	0.28	6,571	1.61	5,364	2.04	1,064	2.15
Q3	8,125	-0.16	6,570	-0.37	5,364	-0.51	1,064	-3.28
Q4	8,125	-0.85	6,571	-2.26	5,365	-4.09	1,065	-12.05
Q1Q4 Diff		1.71		5.85		9.55		24.97





If you are interested to know more about the Angus HeiferSELECT
Technical Specifications please contact Angus Australia:

P: 02 6773 4600 | E: office@angusaustralia.com.au



www.angusaustralia.com.au